

Datenstrukturen und Algorithmen

Exercise 1

FS 2019

Schedule for today


1 Exercise Process

2 Repetition Theory


- Asymptotic Running Time

3 Programming Exercise

Process for the exercises



Process for the exercises



- Thursday:
 - Handout of new exercise sheet (online per Code Expert).
 - Submission of old exercise sheet (online per Code Expert).
- Friday during exercise class:
 - Kick-off presentation of new exercise sheet.
 - Discussion of old exercise sheet.
 - Opportunity to ask questions about lecture and exercises.

2. Repetition Theory

Warm-up

- What is a problem?

Warm-up

- What is a problem?
- What is an algorithm?

Warm-up

- What is a problem?
- What is an algorithm?
 - well-defined computing procedure to compute output data from input data.


Warm-up

- What is a problem?
- What is an algorithm?
 - well-defined computing procedure to compute output data from input data.
- What is a program?


Warm-up

- What is a problem?
- What is an algorithm?
 - well-defined computing procedure to compute output data from input data.
- What is a program?
 - Concrete implementation of an algorithm

Warm-up



Warm-up



Efficiency

Problem	Complexity	Minimal (asymptotic) cost over all algorithms that solve the problem.
Algorithm	Cost	Number of elementary operations
Program	Computing time	Measurable value on an actual machine.

Efficiency

Problem	Complexity	Minimal (asymptotic) cost over all algorithms that solve the problem.
Algorithm	Cost	Number of elementary operations
Program	Computing time	Measurable value on an actual machine.

- Estimation of cost or computing time depending on the input size, denoted by n .

Asymptotic behavior

- What are $\Omega(g(n))$, $\Theta(g(n))$, $\mathcal{O}(g(n))$?

Asymptotic behavior

- What are $\Omega(g(n))$, $\Theta(g(n))$, $\mathcal{O}(g(n))$?
→ Sets of functions!

Asymptotic behavior

- What are $\Omega(g(n))$, $\Theta(g(n))$, $\mathcal{O}(g(n))$?
- Sets of functions!

Repetition, sets A, B :

subset $A \subseteq B$

proper subset $A \subsetneq B$

intersection $A \cap B$

Asymptotic behavior

Given: function $f : \mathbb{N} \rightarrow \mathbb{R}$.

Definition:

$$\mathcal{O}(g) = \{f : \mathbb{N} \rightarrow \mathbb{R} \mid \exists c > 0, n_0 \in \mathbb{N} : 0 \leq f(n) \leq c \cdot g(n) \ \forall n \geq n_0\}$$

$$\Omega(g) = \{f : \mathbb{N} \rightarrow \mathbb{R} \mid \exists c > 0, n_0 \in \mathbb{N} : 0 \leq c \cdot g(n) \leq f(n) \ \forall n \geq n_0\}$$

$$\Theta(g) = \mathcal{O}(g) \cap \Omega(g)$$

Useful information for the exercise

Theorem

- 1 $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0 \Rightarrow f \in \mathcal{O}(g), \mathcal{O}(f) \subsetneq \mathcal{O}(g).$
- 2 $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = C > 0$ (*C constant*) $\Rightarrow f \in \Theta(g).$
- 3 $\frac{f(n)}{g(n)} \xrightarrow[n \rightarrow \infty]{} \infty \Rightarrow g \in \mathcal{O}(f), \mathcal{O}(g) \subsetneq \mathcal{O}(f).$

Useful information for the exercise

Theorem

- 1 $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0 \Rightarrow f \in \mathcal{O}(g), \mathcal{O}(f) \subsetneq \mathcal{O}(g).$
- 2 $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = C > 0$ (*C constant*) $\Rightarrow f \in \Theta(g).$
- 3 $\frac{f(n)}{g(n)} \xrightarrow[n \rightarrow \infty]{} \infty \Rightarrow g \in \mathcal{O}(f), \mathcal{O}(g) \subsetneq \mathcal{O}(f).$

Example

- 1 $\lim_{n \rightarrow \infty} \frac{n}{n^2} = 0 \Rightarrow n \in \mathcal{O}(n^2), \mathcal{O}(n) \subsetneq \mathcal{O}(n^2).$
- 2 $\lim_{n \rightarrow \infty} \frac{2n}{n} = 2 > 0 \Rightarrow 2n \in \Theta(n).$
- 3 $\frac{n^2}{n} \xrightarrow[n \rightarrow \infty]{} \infty \Rightarrow n \in \mathcal{O}(n^2), \mathcal{O}(n) \subsetneq \mathcal{O}(n^2).$

Examples

Examples

- $n \in \mathcal{O}(n^2)$

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong:

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!
- $\mathcal{O}(n) \subseteq \mathcal{O}(n^2)$

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!
- $\mathcal{O}(n) \subseteq \mathcal{O}(n^2)$ is correct

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!
- $\mathcal{O}(n) \subseteq \mathcal{O}(n^2)$ is correct

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!
- $\mathcal{O}(n) \subseteq \mathcal{O}(n^2)$ is correct
- $\Theta(n) \subseteq \Theta(n^2)$

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!
- $\mathcal{O}(n) \subseteq \mathcal{O}(n^2)$ is correct
- $\Theta(n) \subseteq \Theta(n^2)$ is wrong

Examples

- $n \in \mathcal{O}(n^2)$ correct, but too imprecise:
 $n \in \mathcal{O}(n)$ and even $n \in \Theta(n)$.
- $3n^2 \in \mathcal{O}(2n^2)$ correct but uncommon:
Omit constants: $3n^2 \in \mathcal{O}(n^2)$.
- $2n^2 \in \mathcal{O}(n)$ is wrong: $\frac{2n^2}{n} = 2n \xrightarrow{n \rightarrow \infty} \infty$!
- $\mathcal{O}(n) \subseteq \mathcal{O}(n^2)$ is correct
- $\Theta(n) \subseteq \Theta(n^2)$ is wrong $n \notin \Omega(n^2) \supset \Theta(n^2)$

Quiz

$1 \in \mathcal{O}(15)$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$
 $2n + 1 \in \Theta(n)$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$
 $2n + 1 \in \Theta(n)$? ✓

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

$\sqrt{n} \notin \Theta(n)$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

$\sqrt{n} \notin \Theta(n)$? ✓

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

$\sqrt{n} \notin \Theta(n)$? ✓

$\mathcal{O}(\sqrt{n}) \subset \mathcal{O}(n)$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

$\sqrt{n} \notin \Theta(n)$? ✓

$\mathcal{O}(\sqrt{n}) \subset \mathcal{O}(n)$? ✓

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

$\sqrt{n} \notin \Theta(n)$? ✓

$\mathcal{O}(\sqrt{n}) \subset \mathcal{O}(n)$? ✓

$2^n \notin \mathcal{O}(\exp(n))$?

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

$\sqrt{n} \notin \Theta(n)$? ✓

$\mathcal{O}(\sqrt{n}) \subset \mathcal{O}(n)$? ✓

$2^n \notin \mathcal{O}(\exp(n))$? ✗

Quiz

$1 \in \mathcal{O}(15)$? ✓ better $1 \in \mathcal{O}(1)$

$2n + 1 \in \Theta(n)$? ✓

$\sqrt{n} \in \mathcal{O}(n)$? ✓

$\sqrt{n} \in \Omega(n)$? ✗

$n \in \Omega(\sqrt{n})$? ✓

$\sqrt{n} \notin \Theta(n)$? ✓

$\mathcal{O}(\sqrt{n}) \subset \mathcal{O}(n)$? ✓

$2^n \notin \mathcal{O}(\exp(n))$? ✗

Quiz: A good strategy?

... Then I simply buy a new machine

Quiz: A good strategy?

... Then I simply buy a new machine If today I can solve a problem of size n , then with a 10 or 100 times faster machine I can solve ...¹

Komplexität	(speed $\times 10$)	(speed $\times 100$)
$\log_2 n$		
n		
n^2		
2^n		

Quiz: A good strategy?

... Then I simply buy a new machine If today I can solve a problem of size n , then with a 10 or 100 times faster machine I can solve ...¹

Komplexität	(speed $\times 10$)	(speed $\times 100$)
$\log_2 n$	$n \rightarrow n^{10}$	$n \rightarrow n^{100}$
n		
n^2		
2^n		

Quiz: A good strategy?

... Then I simply buy a new machine If today I can solve a problem of size n , then with a 10 or 100 times faster machine I can solve ...¹

Komplexität	(speed $\times 10$)	(speed $\times 100$)
$\log_2 n$	$n \rightarrow n^{10}$	$n \rightarrow n^{100}$
n	$n \rightarrow 10 \cdot n$	$n \rightarrow 100 \cdot n$
n^2		
2^n		

Quiz: A good strategy?

... Then I simply buy a new machine If today I can solve a problem of size n , then with a 10 or 100 times faster machine I can solve ...¹

Komplexität	(speed $\times 10$)	(speed $\times 100$)
$\log_2 n$	$n \rightarrow n^{10}$	$n \rightarrow n^{100}$
n	$n \rightarrow 10 \cdot n$	$n \rightarrow 100 \cdot n$
n^2	$n \rightarrow 3.16 \cdot n$	$n \rightarrow 10 \cdot n$
2^n		

Quiz: A good strategy?

... Then I simply buy a new machine If today I can solve a problem of size n , then with a 10 or 100 times faster machine I can solve ...¹

Komplexität	(speed $\times 10$)	(speed $\times 100$)
$\log_2 n$	$n \rightarrow n^{10}$	$n \rightarrow n^{100}$
n	$n \rightarrow 10 \cdot n$	$n \rightarrow 100 \cdot n$
n^2	$n \rightarrow 3.16 \cdot n$	$n \rightarrow 10 \cdot n$
2^n	$n \rightarrow n + 3.32$	$n \rightarrow n + 6.64$

¹To see this, you set $f(n') = c \cdot f(n)$ ($c = 10$ or $c = 100$) and solve for n'

Asymptotic Running Times with Θ

```
void run(int n){  
    for (int i = 1; i<n; ++i)  
        for (int j = 1; j<n; ++j)  
            op();  
}
```

How often is `op()` called?

Asymptotic Running Times with Θ

```
void run(int n){  
    for (int i = 1; i<n; ++i)  
        for (int j = i; j<n; ++j)  
            op();  
}
```

How often is `op()` called?

Asymptotic Running Times with Θ

```
void run(int n){  
    for (int i = 1; i<n; ++i){  
        op();  
        for (int j = i; j<n; ++j)  
            op();  
    }  
}
```

How often is `op()` called?

Asymptotic Running Times with Θ

```
void run(int n){  
    for(int i = 1; i <= n; ++i)  
        for(int j = 1; j*j <= n; ++j)  
            for(int k = n; k >= 2; --k)  
                op();  
}
```

How often is `op()` called?

3. Programming Exercise

Sum in sub-interval (naive algorithm)

Input: A sequence of n numbers $(a_0, a_1, \dots, a_{n-1})$ and a sub-interval

$$I = [x_0, x_1]$$

Output: $\sum_{i=x_0}^{x_1} a_i.$

$S \leftarrow 0$

for $i \in \{x_0, \dots, x_1\}$ **do**

$S \leftarrow S + a_i$

return S

Sum in sub-interval (naive algorithm)

Input: A sequence of n numbers $(a_0, a_1, \dots, a_{n-1})$ and a sub-interval

$$I = [x_0, x_1]$$

Output: $\sum_{i=x_0}^{x_1} a_i.$

$S \leftarrow 0$

for $i \in \{x_0, \dots, x_1\}$ **do**

$S \leftarrow S + a_i$

return S

Idea

- Use the prefix sum to compute the sum of arbitrary sub-intervals with constant complexity
- Generalization to two dimensions.

Multidimensional vectors

Definition

```
std::vector< std::vector<int> > my_vec( n_rows,  
std::vector<int>(n_cols,init_value) );
```

Indexing

```
my_vec[row] [col]
```

Classes

```
class Insurance { // Definition
public: // public section
    Insurance(double rate) {rate_ = rate;} // Konstruktor
    double get_rate() {return rate_;} // member function
private: // private section
    double rate_; // data member
};

int main() {
    Insurance insurance(2.);
    std::cout << insurance.get_rate();
    return 0;
}
```

Questions or Suggestions?